

Phenomenology of warped extra dimensions at LHC

V. Ravindran

Harish-Chandra Research Institute, Allahabad

- TeV scale gravity models
- Phenomenology with them
- NLO QCD results for Drell-Yan at TeV scale
- Summary

In collaboration with

Prakash Mathews, K. Sridhar, W. L. van Neerven & M.C. Kumar

Large Extra Dimensions

- Models of "Extra Dimensions" are now studied as serious contenders for "Physics Beyond SM"(BSM).
- They provide an alternate view of the "hierarchy" between the EW ($\sim 1 \text{ TeV}$) and the Planck scale (10^{16} TeV)

Large Extra Dimensions

- Models of "Extra Dimensions" are now studied as serious contenders for "Physics Beyond SM"(BSM).
- They provide an alternate view of the "hierarchy" between the EW ($\sim 1 \text{ TeV}$) and the Planck scale (10^{16} TeV)
- The "Geometry" of extra spacial dimensions may be responsible for the "Hierarchy"

Large vol of extra dim ADD ADD PLB 429(1998)	Strong Curvature of extra dim RS RS PRL 83 (1999) 3370
--	--

Large Extra Dimensions

- Models of "Extra Dimensions" are now studied as serious contenders for "Physics Beyond SM"(BSM).
- They provide an alternate view of the "hierarchy" between the EW ($\sim 1 \text{ TeV}$) and the Planck scale (10^{16} TeV)
- The "Geometry" of extra spacial dimensions may be responsible for the "Hierarchy"

Large vol of extra dim ADD ADD PLB 429(1998)	Strong Curvature of extra dim RS RS PRL 83 (1999) 3370
--	--


- A "viable mechanism" to "hide" the "extra dim" such that "space time" is effectively "four" consistent with "known physics".

Large Extra Dimensions

- Models of "Extra Dimensions" are now studied as serious contenders for "Physics Beyond SM"(BSM).
- They provide an alternate view of the "hierarchy" between the EW ($\sim 1 \text{ TeV}$) and the Planck scale (10^{16} TeV)
- The "Geometry" of extra spacial dimensions may be responsible for the "Hierarchy"

Large vol of extra dim ADD ADD PLB 429(1998)	Strong Curvature of extra dim RS RS PRL 83 (1999) 3370
--	--

- A "viable mechanism" to "hide" the "extra dim" such that "space time" is effectively "four" consistent with "known physics".
- No deviation at present accelerators leads to limits on extra dim $< 10^{-16} \text{ cm}$




Large Extra Dimensions

- Models of "Extra Dimensions" are now studied as serious contenders for "Physics Beyond SM"(BSM).
- They provide an alternate view of the "hierarchy" between the EW ($\sim 1 \text{ TeV}$) and the Planck scale (10^{16} TeV)
- The "Geometry" of extra spacial dimensions may be responsible for the "Hierarchy"

Large vol of extra dim ADD ADD PLB 429(1998)	Strong Curvature of extra dim RS RS PRL 83 (1999) 3370
--	--

- A "viable mechanism" to "hide" the "extra dim" such that "space time" is effectively "four" consistent with "known physics".
- No deviation at present accelerators leads to limits on extra dim $< 10^{-16} \text{ cm}$



Large Extra Dimensions (ADD Scenario)

- Space time $\mathcal{M}_4 \times \mathcal{K}_d$ FACTORISABLE GEOMETRY


\mathcal{M}_4 : 3+1 dim space time \mathcal{K}_d : Compact space of size R

Large Extra Dimensions (ADD Scenario)

- Space time $\mathcal{M}_4 \times \mathcal{K}_d$ FACTORISABLE GEOMETRY

\mathcal{M}_4 : 3+1 dim space time \mathcal{K}_d : Compact space of size R

- SM localised on a 3-brane embedded into the 4+d dim space time with d compact extra dimensions which can ONLY be probed by gravity




Large Extra Dimensions (ADD Scenario)

- Space time $\mathcal{M}_4 \times \mathcal{K}_d$ FACTORISABLE GEOMETRY

\mathcal{M}_4 : 3+1 dim space time \mathcal{K}_d : Compact space of size R

- SM localised on a 3-brane embedded into the 4+d dim space time with d compact extra dimensions which can ONLY be probed by gravity



- Gauss law in 4+d dim with d compact dim of radius R

$$r \ll R$$

$$r \gg R$$

$$V(r) \sim \frac{m_1 m_2}{M_S^{d+2}} \frac{1}{r^{d+1}}$$

$$V(r) \sim \frac{m_1 m_2}{M_S^{d+2} R^d} \frac{1}{r}$$

Large Extra Dimensions

- $r \gg R$ gravitational flux lines in 4+d dim are constrained in the compact dim and hence the potential is effectively r^{-1} at large distances

$$\boxed{M_P^2 \sim M_S^{2+d} R^d}$$

- M_P : 4 dim Planck scale
- M_S : $\sim \text{TeV}$ 4+d dim Planck scale
- R : radius of compactification, could be large compared to a TeV^{-1}

Large Extra Dimensions

- $r \gg R$ gravitational flux lines in 4+d dim are constrained in the compact dim and hence the potential is effectively r^{-1} at large distances

$$\boxed{M_P^2 \sim M_S^{2+d} R^d}$$

- M_P : 4 dim Planck scale
- M_S : $\sim \text{TeV}$ 4+d dim Planck scale
- R : radius of compactification, could be large compared to a TeV^{-1}

$$R \sim 10^{\frac{30}{d}-17} \text{ cm}$$

$$R^{-1} \sim 10^{\frac{-30}{d}+3} \text{ GeV}$$

d	$R \text{ (cm)}$	R^{-1}
1	10^{13}	10^{-27} eV
2	10^{-2}	10^{-3} eV
3	10^{-7}	100 eV
...
6	10^{-12}	10 MeV

Large Extra Dimensions

- $r \gg R$ gravitational flux lines in 4+d dim are constrained in the compact dim and hence the potential is effectively r^{-1} at large distances

$$\boxed{M_P^2 \sim M_S^{2+d} R^d}$$

- M_P : 4 dim Planck scale
- M_S : $\sim \text{TeV}$ 4+d dim Planck scale
- R : radius of compactification, could be large compared to a TeV^{-1}

$$R \sim 10^{\frac{30}{d}-17} \text{ cm}$$

$$R^{-1} \sim 10^{\frac{-30}{d}+3} \text{ GeV}$$

d	$R \text{ (cm)}$	R^{-1}
1	10^{13}	10^{-27} eV
2	10^{-2}	10^{-3} eV
3	10^{-7}	100 eV
...
6	10^{-12}	10 MeV

- Only Gravitational field can probe the full 4+n dim space, deviation from Newtonian gravity puts constraint on number of extra dim

$$d \geq 2 \text{ Possible}$$


If Fundamental Planck Scale $M_S \sim 1 \text{ TeV}$, "no Hierarchy problem".

Kaluza-Klein Modes

- Extra dimensions being compact, gravitational field will be periodic function in the extra dimension.
- In 4-dim it would correspond to nearly mass degenerate tower of KK modes $m_{\vec{n}}^2 \sim \vec{n}^2 / R^2$


Kaluza-Klein Modes

- Extra dimensions being compact, gravitational field will be periodic function in the extra dimension.
- In 4-dim it would correspond to nearly mass degenerate tower of KK modes $m_{\vec{n}}^2 \sim \vec{n}^2 / R^2$



Massless graviton and KK modes couple with SM fields with coupling $M_P^{-1} \sim R^{\frac{d}{2}}$


Kaluza-Klein suppression



The diagram shows an annihilation process where a fermion q and an antifermion \bar{q} meet at a vertex. A dashed line representing a Kaluza-Klein gauge boson, labeled $\sum_n G^{\vec{n}}$, connects this vertex to another vertex where a fermion and an antifermion are produced.

$$\sim \frac{1}{R^{\frac{d}{2}}} \sum_n \frac{1}{Q^2 - m_{\vec{n}}^2 + i\epsilon} \frac{1}{R^{\frac{d}{2}}}$$

Kaluza-Klein suppression




A Feynman diagram illustrating the annihilation of a fermion q and an antifermion \bar{q} into a scalar particle. The incoming fermion q and antifermion \bar{q} lines meet at a vertex, and the outgoing scalar particle is represented by a dashed line. The interaction is labeled $\sum_n G^{\vec{n}}$. To the right of the diagram, the expression is given as:

$$\sim \frac{1}{R^{\frac{d}{2}}} \sum_n \frac{1}{Q^2 - m_{\vec{n}}^2 + i\epsilon} \frac{1}{R^{\frac{d}{2}}}$$

- State density of KK modes $\sim V_d$

$$\rho(m_{\vec{n}}) = \frac{R^d m_{\vec{n}}^{d-2}}{(4\pi)^{d/2} \Gamma(d/2)}$$

Kaluza-Klein suppression



$$\sim \frac{1}{R^{\frac{d}{2}}} \sum_n \frac{1}{Q^2 - m_{\vec{n}}^2 + i\epsilon} \frac{1}{R^{\frac{d}{2}}}$$


- State density of KK modes $\sim V_d$

$$\rho(m_{\vec{n}}) = \frac{R^d m_{\vec{n}}^{d-2}}{(4\pi)^{d/2} \Gamma(d/2)}$$

- (Coupling)² (Sum over the KK mode) leads to

$$\left(\frac{1}{R^{\frac{d}{2}}} \right)^2 \sum_n \frac{1}{Q^2 - m_{\vec{n}}^2 + i\epsilon} \Rightarrow \frac{1}{R^d} V_d \sim \frac{1}{R^d} \quad C(d, M_S) \quad R^d \Rightarrow \frac{1}{M_S^{4+d}}$$

Kaluza-Klein suppression



$$\sim \frac{1}{R^{\frac{d}{2}}} \sum_n \frac{1}{Q^2 - m_{\vec{n}}^2 + i\epsilon} \frac{1}{R^{\frac{d}{2}}}$$

- State density of KK modes $\sim V_d$

$$\rho(m_{\vec{n}}) = \frac{R^d m_{\vec{n}}^{d-2}}{(4\pi)^{d/2} \Gamma(d/2)}$$

- (Coupling)² (Sum over the KK mode) leads to

$$\left(\frac{1}{R^{\frac{d}{2}}}\right)^2 \sum_n \frac{1}{Q^2 - m_{\vec{n}}^2 + i\epsilon} \Rightarrow \frac{1}{R^d} V_d \sim \frac{1}{R^d} \quad C(d, M_S) \quad R^d \Rightarrow \frac{1}{M_S^{4+d}}$$

$$\frac{1}{M_S^{4+d}} \sim \frac{1}{(TeV)^{4+d}}$$

Planck suppression is compensated by High multiplicity of KK modes

Kaluza-Klein suppression

$$\left| \begin{array}{c} q \\ \bar{q} \end{array} \right\} \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} \text{---} \xrightarrow{\sum_n G^{\vec{n}}} \text{---} + \dots \end{array} \left| \right|^2 \sim \left(\frac{1}{R^{\frac{d}{2}}} \right)^2 \int dP S_{\vec{n}} \delta^+(s - m_{\vec{n}}^2)$$

Kaluza-Klein suppression

$$\left| \begin{array}{c} q \\ \bar{q} \end{array} \right\} \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} \sum_n G^{\vec{n}} \\ \text{---} \rightarrow \text{---} + \dots \end{array} \left| \right|^2 \sim \left(\frac{1}{R^{\frac{d}{2}}} \right)^2 \int dP S_{\vec{n}} \delta^+(s - m_{\vec{n}}^2)$$

- State density of KK modes $\sim V_d$

$$\rho(m_{\vec{n}}) = \frac{R^d m_{\vec{n}}^{d-2}}{(4\pi)^{d/2} \Gamma(d/2)}$$

Kaluza-Klein suppression

$$\left| \begin{array}{c} q \\ \bar{q} \end{array} \right\} \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} \sum_n G^{\vec{n}} \\ \text{---} \rightarrow \text{---} + \dots \end{array} \left| \right|^2 \sim \left(\frac{1}{R^{\frac{d}{2}}} \right)^2 \int dP S_{\vec{n}} \delta^+(s - m_{\vec{n}}^2)$$

- State density of KK modes $\sim V_d$

$$\rho(m_{\vec{n}}) = \frac{R^d m_{\vec{n}}^{d-2}}{(4\pi)^{d/2} \Gamma(d/2)}$$

- (Coupling)² (Sum over the KK mode) leads to

$$\left(\frac{1}{R^{\frac{d}{2}}} \right)^2 \int dP S_{\vec{n}} \delta^+(s - m_{\vec{n}}^2) \Rightarrow \frac{1}{R^d} V_d \sim \frac{1}{R^d} C'(d, M_S) R^d \Rightarrow \frac{1}{M_S^{4+d}}$$

Kaluza-Klein suppression

$$\left| \begin{array}{c} q \\ \bar{q} \end{array} \right\rangle \left[\begin{array}{c} \sum_n G^{\vec{n}} \\ \text{---} \end{array} \right] \left| \right|^2 \sim \left(\frac{1}{R^{\frac{d}{2}}} \right)^2 \int dP S_{\vec{n}} \delta^+(s - m_{\vec{n}}^2)$$

- State density of KK modes $\sim V_d$

$$\rho(m_{\vec{n}}) = \frac{R^d m_{\vec{n}}^{d-2}}{(4\pi)^{d/2} \Gamma(d/2)}$$

- (Coupling)² (Sum over the KK mode) leads to

$$\left(\frac{1}{R^{\frac{d}{2}}} \right)^2 \int dP S_{\vec{n}} \delta^+(s - m_{\vec{n}}^2) \Rightarrow \frac{1}{R^d} V_d \sim \frac{1}{\mathbb{R}^d} C'(d, M_S) \mathbb{R}^d \Rightarrow \frac{1}{M_S^{4+d}}$$

$$\frac{1}{M_S^{4+d}} \sim \frac{1}{(TeV)^{4+d}}$$

Planck suppression is compensated by High multiplicity of KK modes

Warped Extra Dimension

Warped Extra Dimension

- Non-Factorisable geometry, 5-dim AdS space— constant negative curvature

$$ds^2 = e^{-2\mathcal{K}r_c|\phi|} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2$$

Warped Extra Dimension

- Non-Factorisable geometry, 5-dim AdS space— constant negative curvature

$$ds^2 = e^{-2\mathcal{K}r_c|\phi|} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2$$

- Extra 5th dim compactified to the orbifold S^1/Z_2

$$S^1 : y \equiv y + 2\pi R \quad \text{and} \quad Z_2 : y \equiv -y$$

leads to compactification on a segment of size πR .

Warped Extra Dimension

- Non-Factorisable geometry, 5-dim AdS space— constant negative curvature

$$ds^2 = e^{-2\mathcal{K}r_c|\phi|} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2$$

- Extra 5th dim compactified to the orbifold S^1/Z_2

$$S^1 : y \equiv y + 2\pi R \quad \text{and} \quad Z_2 : y \equiv -y$$

leads to compactification on a segment of size πR .

- Two 3-Branes located at orbifold fixed points $[0, \pi R]$.

Warped Extra Dimension

- Non-Factorisable geometry, 5-dim AdS space— constant negative curvature

$$ds^2 = e^{-2\mathcal{K}r_c|\phi|} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2$$

- Extra 5th dim compactified to the orbifold S^1/Z_2

$$S^1 : y \equiv y + 2\pi R \quad \text{and} \quad Z_2 : y \equiv -y$$

leads to compactification on a segment of size πR .

- Two 3-Branes located at orbifold fixed points $[0, \pi R]$.
- TeV Brane (SM constrained) at $y = \pi R$

Warped Extra Dimension

- Non-Factorisable geometry, 5-dim AdS space— constant negative curvature

$$ds^2 = e^{-2\mathcal{K}r_c|\phi|} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2$$

- Extra 5th dim compactified to the orbifold S^1/Z_2

$$S^1 : y \equiv y + 2\pi R \quad \text{and} \quad Z_2 : y \equiv -y$$

leads to compactification on a segment of size πR .

- Two 3-Branes located at orbifold fixed points $[0, \pi R]$.
- TeV Brane (SM constrained) at $y = \pi R$
- Planck Brane (Gravity resides) at $y = 0$

Warped Extra Dimension

- Non-Factorisable geometry, 5-dim AdS space— constant negative curvature

$$ds^2 = e^{-2\mathcal{K}r_c|\phi|} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2$$

- Extra 5th dim compactified to the orbifold S^1/Z_2

$$S^1 : y \equiv y + 2\pi R \quad \text{and} \quad Z_2 : y \equiv -y$$

leads to compactification on a segment of size πR .

- Two 3-Branes located at orbifold fixed points $[0, \pi R]$.
- TeV Brane (SM constrained) at $y = \pi R$
- Planck Brane (Gravity resides) at $y = 0$
- gravity propagates the bulk.

Warped Extra Dimension

Warped Extra Dimension

- Interaction of RS KK tower with SM fields on the TEV Brane

$$\mathcal{L} \sim -\frac{1}{M_P} T^{\alpha\beta} G_{\alpha\beta}^{(0)} - \frac{1}{\Lambda_\pi} T^{\alpha\beta} \sum_{n=1}^{\infty} G_{\alpha\beta}^{(n)}$$

$$\Lambda_\pi \sim M_P e^{-\mathcal{K} r_c \pi} \sim \mathcal{O}(\text{TeV})$$

Warped Extra Dimension

- Interaction of RS KK tower with SM fields on the TEV Brane

$$\mathcal{L} \sim -\frac{1}{M_P} T^{\alpha\beta} G_{\alpha\beta}^{(0)} - \frac{1}{\Lambda_\pi} T^{\alpha\beta} \sum_{n=1}^{\infty} G_{\alpha\beta}^{(n)}$$

$$\Lambda_\pi \sim M_P e^{-\mathcal{K} r_c \pi} \sim \mathcal{O}(\text{TeV})$$

- Zero mode decouples (massless graviton) Newtonian Gravity intact M_P^{-1}

Warped Extra Dimension

- Interaction of RS KK tower with SM fields on the TEV Brane

$$\mathcal{L} \sim -\frac{1}{M_P} T^{\alpha\beta} G_{\alpha\beta}^{(0)} - \frac{1}{\Lambda_\pi} T^{\alpha\beta} \sum_{n=1}^{\infty} G_{\alpha\beta}^{(n)}$$

$$\Lambda_\pi \sim M_P e^{-\mathcal{K} r_c \pi} \sim \mathcal{O}(\text{TeV})$$

- Zero mode decouples (massless graviton) Newtonian Gravity intact M_P^{-1}
- Excited massive KK modes couple to SM with TeV^{-1} suppression

$$M_n = x_n \mathcal{K} e^{-\mathcal{K} r_c \pi} \equiv x_n m_0$$

Warped Extra Dimension

- Interaction of RS KK tower with SM fields on the TEV Brane

$$\mathcal{L} \sim -\frac{1}{M_P} T^{\alpha\beta} G_{\alpha\beta}^{(0)} - \frac{1}{\Lambda_\pi} T^{\alpha\beta} \sum_{n=1}^{\infty} G_{\alpha\beta}^{(n)}$$

$$\Lambda_\pi \sim M_P e^{-\mathcal{K} r_c \pi} \sim \mathcal{O}(\text{TeV})$$

- Zero mode decouples (massless graviton) Newtonian Gravity intact M_P^{-1}
- Excited massive KK modes couple to SM with TeV^{-1} suppression

$$M_n = x_n \mathcal{K} e^{-\mathcal{K} r_c \pi} \equiv x_n m_0$$

- x_n - the zeros of the Bessel function $J_1(x)$
- The zero-mode couples weakly and decouples
- The couplings of the massive RS gravitons are enhanced by the exponential $e^{\pi \mathcal{K} R_c}$ leading to interactions of electroweak strength.

Warped Extra Dimension

- Interaction of RS KK tower with SM fields on the TEV Brane

$$\mathcal{L} \sim -\frac{1}{M_P} T^{\alpha\beta} G_{\alpha\beta}^{(0)} - \frac{1}{\Lambda_\pi} T^{\alpha\beta} \sum_{n=1}^{\infty} G_{\alpha\beta}^{(n)}$$

$$\Lambda_\pi \sim M_P e^{-\mathcal{K} r_c \pi} \sim \mathcal{O}(\text{TeV})$$

- Zero mode decouples (massless graviton) Newtonian Gravity intact M_P^{-1}
- Excited massive KK modes couple to SM with TeV^{-1} suppression

$$M_n = x_n \mathcal{K} e^{-\mathcal{K} r_c \pi} \equiv x_n m_0$$

- x_n - the zeros of the Bessel function $J_1(x)$
- The zero-mode couples weakly and decouples
- The couplings of the massive RS gravitons are enhanced by the exponential $e^{\pi \mathcal{K} R_c}$ leading to interactions of electroweak strength.
- Two basic parameters of the RS model are

$$m_0 = \mathcal{K} e^{-\mathcal{K} r_c \pi} \quad c_0 = \frac{\mathcal{K}}{M_P}$$

Warped Extra Dimension

Warped Extra Dimension

The propagator:

$$P_G(q) = \mathcal{D}(Q^2) B_{\mu\nu\lambda\rho}(q) ,$$

$$B_{\mu\nu\rho\sigma}(q) = \left(g_{\mu\rho} - \frac{q_\mu q_\rho}{M_n^2} \right) \left(g_{\nu\sigma} - \frac{q_\nu q_\sigma}{M_n^2} \right) + \left(g_{\mu\sigma} - \frac{q_\mu q_\sigma}{M_n^2} \right) \left(g_{\nu\rho} - \frac{q_\nu q_\rho}{M_n^2} \right) \\ - \frac{2}{n-1} \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{M_n^2} \right) \left(g_{\rho\sigma} - \frac{q_\rho q_\sigma}{M_n^2} \right) .$$

Warped Extra Dimension

The propagator:

$$P_G(q) = \mathcal{D}(Q^2) B_{\mu\nu\lambda\rho}(q) ,$$

$$B_{\mu\nu\rho\sigma}(q) = \left(g_{\mu\rho} - \frac{q_\mu q_\rho}{M_n^2} \right) \left(g_{\nu\sigma} - \frac{q_\nu q_\sigma}{M_n^2} \right) + \left(g_{\mu\sigma} - \frac{q_\mu q_\sigma}{M_n^2} \right) \left(g_{\nu\rho} - \frac{q_\nu q_\rho}{M_n^2} \right) \\ - \frac{2}{n-1} \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{M_n^2} \right) \left(g_{\rho\sigma} - \frac{q_\rho q_\sigma}{M_n^2} \right) .$$

$$\mathcal{D}(Q^2) = \sum_{n=1}^{\infty} \frac{1}{Q^2 - M_n^2 + iM_n\Gamma_n} \equiv \frac{\lambda}{m_0^2} ,$$

M_n -the masses of the individual resonances, Γ_n are the widths.

Warped Extra Dimension

The propagator:

$$P_G(q) = \mathcal{D}(Q^2) B_{\mu\nu\lambda\rho}(q) ,$$

$$B_{\mu\nu\rho\sigma}(q) = \left(g_{\mu\rho} - \frac{q_\mu q_\rho}{M_n^2} \right) \left(g_{\nu\sigma} - \frac{q_\nu q_\sigma}{M_n^2} \right) + \left(g_{\mu\sigma} - \frac{q_\mu q_\sigma}{M_n^2} \right) \left(g_{\nu\rho} - \frac{q_\nu q_\rho}{M_n^2} \right) \\ - \frac{2}{n-1} \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{M_n^2} \right) \left(g_{\rho\sigma} - \frac{q_\rho q_\sigma}{M_n^2} \right) .$$

$$\mathcal{D}(Q^2) = \sum_{n=1}^{\infty} \frac{1}{Q^2 - M_n^2 + iM_n\Gamma_n} \equiv \frac{\lambda}{m_0^2} ,$$

M_n -the masses of the individual resonances, Γ_n are the widths. λ is defined as

$$\lambda(x_s) = \sum_{n=1}^{\infty} \frac{x_s^2 - x_n^2 - i\frac{\Gamma_n}{m_0}x_n}{x_s^2 - x_n^2 + \frac{\Gamma_n}{m_0}x_n} , \quad x_s = Q/m_0 .$$

We have to sum over all the resonances to get the value of $\lambda(x_s)$.

Phenomenology with Extra-Dimension

In the SM, the partonic cross sections decreases with the energy scale (Q or p_T involved):

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{SM}(\hat{s}, Q^2) \sim \frac{1}{\hat{s} Q^2}$$

Phenomenology with Extra-Dimension

In the SM, the partonic cross sections decreases with the energy scale (Q or p_T involved):

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{SM}(\hat{s}, Q^2) \sim \frac{1}{\hat{s} Q^2}$$

In the ADD, the partonic cross sections increase monotonically with the energy scale involved:

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{ADD}(\hat{s}, Q^2, M_S) \sim \left(\frac{Q^2}{M_S^2} \right)^{d-2} \frac{Q^6}{\hat{s} M_S^8}, \quad Q < M_S$$

Phenomenology with Extra-Dimension

In the SM, the partonic cross sections decreases with the energy scale (Q or p_T involved):

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{SM}(\hat{s}, Q^2) \sim \frac{1}{\hat{s} Q^2}$$

In the ADD, the partonic cross sections increase monotonically with the energy scale involved:

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{ADD}(\hat{s}, Q^2, M_S) \sim \left(\frac{Q^2}{M_S^2} \right)^{d-2} \frac{Q^6}{\hat{s} M_S^8}, \quad Q < M_S$$

- Gravity mediated cross sections can show up at high Q .

Phenomenology with Extra-Dimension

In the SM, the partonic cross sections decreases with the energy scale (Q or p_T involved):

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{SM}(\hat{s}, Q^2) \sim \frac{1}{\hat{s} Q^2}$$

In the ADD, the partonic cross sections increase monotonically with the energy scale involved:

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{ADD}(\hat{s}, Q^2, M_S) \sim \left(\frac{Q^2}{M_S^2} \right)^{d-2} \frac{Q^6}{\hat{s} M_S^8}, \quad Q < M_S$$

- Gravity mediated cross sections can show up at high Q .

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{RS}(\hat{s}, Q^2, M_S) \sim c_o^4 \lambda^2 \left(\frac{Q^6}{\hat{s} m_o^8} \right), \quad Q < m_o$$

- The processes where the virtual/real KK gravitons contribute significantly:

- (1) Di-lepton or Drell-Yan production at large invariant mass Q
- (2) Di-photon or Di-boson production at large Q, P_T
- (3) Observables with missing energy (...) . . .

Drell-Yan Process


$$P_1(p_1) + P_2(p_2) \rightarrow [\gamma, Z, \textcolor{red}{G}] + \text{hadronic states}(X)$$

$$\hookrightarrow l^+(k_1) + l^-(k_2) \quad (k_1 + k_2)^2 = Q^2$$

Drell-Yan Process

$$P_1(p_1) + P_2(p_2) \rightarrow [\gamma, Z, \textcolor{red}{G}] + \text{hadronic states}(X)$$



$$\hookrightarrow l^+(k_1) + l^-(k_2) \quad (k_1 + k_2)^2 = Q^2$$



Drell-Yan Process

$$P_1(p_1) + P_2(p_2) \rightarrow [\gamma, Z, \textcolor{red}{G}] + \text{hadronic states}(X)$$

$$\hookrightarrow l^+(k_1) + l^-(k_2) \quad (k_1 + k_2)^2 = Q^2$$





SM


Drell-Yan Process

$$P_1(p_1) + P_2(p_2) \rightarrow [\gamma, Z, \textcolor{red}{G}] + \text{hadronic states}(X)$$

$$\hookrightarrow l^+(k_1) + l^-(k_2) \quad (k_1 + k_2)^2 = Q^2$$





SM





Drell-Yan Process

$$P_1(p_1) + P_2(p_2) \rightarrow [\gamma, Z, \textcolor{red}{G}] + \text{hadronic states}(X)$$

$$\hookrightarrow l^+(k_1) + l^-(k_2) \quad (k_1 + k_2)^2 = Q^2$$



SM



Gravity

Parton Model

Hadronic cross section in terms of partonic cross sections convoluted with appropriate PDF:

$$2S \, d\sigma^{P_1 P_2}(\tau, Q^2) = \sum_{ab} \int_{\tau}^1 \frac{dx}{x} \Phi_{ab}(x, \mu_F) \, 2\hat{s} \, d\hat{\sigma}^{ab}\left(\frac{\tau}{x}, Q^2, \mu_F\right)$$

Parton Model

Hadronic cross section in terms of partonic cross sections convoluted with appropriate PDF:

$$2S d\sigma^{P_1 P_2}(\tau, Q^2) = \sum_{ab} \int_{\tau}^1 \frac{dx}{x} \Phi_{ab}(x, \mu_F) 2\hat{s} d\hat{\sigma}^{ab}\left(\frac{\tau}{x}, Q^2, \mu_F\right)$$

- The perturbatively calculable partonic cross section:

$$d\hat{\sigma}^{ab}(z, Q^2, \mu_F) = \sum_{i=0}^{\infty} \left(\frac{\alpha_s(\mu_R^2)}{4\pi} \right)^i d\hat{\sigma}^{ab,(i)}(z, Q^2, \mu_F, \mu_R) \quad (1)$$

Parton Model

Hadronic cross section in terms of partonic cross sections convoluted with appropriate PDF:

$$2S d\sigma^{P_1 P_2}(\tau, Q^2) = \sum_{ab} \int_{\tau}^1 \frac{dx}{x} \Phi_{ab}(x, \mu_F) 2\hat{s} d\hat{\sigma}^{ab}\left(\frac{\tau}{x}, Q^2, \mu_F\right)$$

- The perturbatively calculable partonic cross section:

$$d\hat{\sigma}^{ab}(z, Q^2, \mu_F) = \sum_{i=0}^{\infty} \left(\frac{\alpha_s(\mu_R^2)}{4\pi} \right)^i d\hat{\sigma}^{ab,(i)}(z, Q^2, \mu_F, \mu_R) \quad (1)$$

- The non-perturbative flux:

$$\Phi_{ab}(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a(z, \mu_F) f_b\left(\frac{x}{z}, \mu_F\right)$$

Parton Model

Hadronic cross section in terms of partonic cross sections convoluted with appropriate PDF:

$$2S d\sigma^{P_1 P_2}(\tau, Q^2) = \sum_{ab} \int_{\tau}^1 \frac{dx}{x} \Phi_{ab}(x, \mu_F) 2\hat{s} d\hat{\sigma}^{ab}\left(\frac{\tau}{x}, Q^2, \mu_F\right)$$

- The perturbatively calculable partonic cross section:

$$d\hat{\sigma}^{ab}(z, Q^2, \mu_F) = \sum_{i=0}^{\infty} \left(\frac{\alpha_s(\mu_R^2)}{4\pi} \right)^i d\hat{\sigma}^{ab,(i)}(z, Q^2, \mu_F, \mu_R) \quad (1)$$

- The non-perturbative flux:

$$\Phi_{ab}(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a(z, \mu_F) f_b\left(\frac{x}{z}, \mu_F\right)$$

- $f_a^{P_1}(x, \mu_F)$ are Parton distribution functions:
- x is the partonic momentum fraction

Parton Model

Hadronic cross section in terms of partonic cross sections convoluted with appropriate PDF:

$$2S d\sigma^{P_1 P_2}(\tau, Q^2) = \sum_{ab} \int_{\tau}^1 \frac{dx}{x} \Phi_{ab}(x, \mu_F) 2\hat{s} d\hat{\sigma}^{ab}\left(\frac{\tau}{x}, Q^2, \mu_F\right)$$

- The perturbatively calculable partonic cross section:

$$d\hat{\sigma}^{ab}(z, Q^2, \mu_F) = \sum_{i=0}^{\infty} \left(\frac{\alpha_s(\mu_R^2)}{4\pi} \right)^i d\hat{\sigma}^{ab,(i)}(z, Q^2, \mu_F, \mu_R) \quad (1)$$

- The non-perturbative flux:

$$\Phi_{ab}(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a(z, \mu_F) f_b\left(\frac{x}{z}, \mu_F\right)$$

- $f_a^{P_1}(x, \mu_F)$ are Parton distribution functions:
- x is the partonic momentum fraction
- μ_R is the Renormalisation scale
- μ_F is the Factorisation scale

Source of Theoretical Uncertainties

Source of Theoretical Uncertainties

- Renormalisation scale
Due to UV divergence at beyond Leading Order

$$\alpha_s \rightarrow \alpha_s(\mu_R^2)$$

Source of Theoretical Uncertainties

- Renormalisation scale
Due to UV divergence at beyond Leading Order

$$\alpha_s \rightarrow \alpha_s(\mu_R^2)$$

- Factorisation scale
Originate from light quarks and massless gluon. Parton distribution functions are renormalised at the factorisation scale μ_F

$$f_a(x) \rightarrow f_a(x, \mu_F^2) \quad a = q, \bar{q}, g$$

Source of Theoretical Uncertainties

- Renormalisation scale
Due to UV divergence at beyond Leading Order

$$\alpha_s \rightarrow \alpha_s(\mu_R^2)$$

- Factorisation scale
Originate from light quarks and massless gluon. Parton distribution functions are renormalised at the factorisation scale μ_F

$$f_a(x) \rightarrow f_a(x, \mu_F^2) \quad a = q, \bar{q}, g$$

- Parton Distribution Functions
Not calculable but extracted from experiments in some factorisation scheme

Source of Theoretical Uncertainties

- Renormalisation scale
Due to UV divergence at beyond Leading Order

$$\alpha_s \rightarrow \alpha_s(\mu_R^2)$$

- Factorisation scale
Originate from light quarks and massless gluon. Parton distribution functions are renormalised at the factorisation scale μ_F

$$f_a(x) \rightarrow f_a(x, \mu_F^2) \quad a = q, \bar{q}, g$$

- Parton Distribution Functions
Not calculable but extracted from experiments in some factorisation scheme
- Observables are "free" of μ_R and μ_F .

Source of Theoretical Uncertainties

- Renormalisation scale
Due to UV divergence at beyond Leading Order

$$\alpha_s \rightarrow \alpha_s(\mu_R^2)$$

- Factorisation scale
Originate from light quarks and massless gluon. Parton distribution functions are renormalised at the factorisation scale μ_F

$$f_a(x) \rightarrow f_a(x, \mu_F^2) \quad a = q, \bar{q}, g$$

- Parton Distribution Functions
Not calculable but extracted from experiments in some factorisation scheme
- Observables are "free" of μ_R and μ_F .
- "Fixed order" perturbative results depend on μ_R and μ_F .

Source of Theoretical Uncertainties

- Renormalisation scale
Due to UV divergence at beyond Leading Order

$$\alpha_s \rightarrow \alpha_s(\mu_R^2)$$

- Factorisation scale
Originate from light quarks and massless gluon. Parton distribution functions are renormalised at the factorisation scale μ_F

$$f_a(x) \rightarrow f_a(x, \mu_F^2) \quad a = q, \bar{q}, g$$

- Parton Distribution Functions
Not calculable but extracted from experiments in some factorisation scheme
- Observables are "free" of μ_R and μ_F .
- "Fixed order" perturbative results depend on μ_R and μ_F .
- Can in principle give large uncertainties.

Source of Theoretical Uncertainties

- Renormalisation scale
Due to UV divergence at beyond Leading Order

$$\alpha_s \rightarrow \alpha_s(\mu_R^2)$$

- Factorisation scale
Originate from light quarks and massless gluon. Parton distribution functions are renormalised at the factorisation scale μ_F



$$f_a(x) \rightarrow f_a(x, \mu_F^2) \quad a = q, \bar{q}, g$$

- Parton Distribution Functions
Not calculable but extracted from experiments in some factorisation scheme
- Observables are "free" of μ_R and μ_F .
- "Fixed order" perturbative results depend on μ_R and μ_F .
- Can in principle give large uncertainties.

It is hence important for extra dimension searches to have better control over the theoretical uncertainties

CTEQ/MRST

Various groups perform global analysis of wide range of DIS and other hadron scattering data to get best fits to a particular order in QCD




○ PDFs satisfies the general constraint, but could differ from each other as experimental and theoretical uncertainties are involved, various assumptions and initial conditions are used

Contributing Subprocess

Leading Order:

Standard Model	Gravity
$q + \bar{q} \rightarrow \gamma/Z$	$q + \bar{q} \rightarrow G$ $g + g \rightarrow G$



Born contributions

Natural question

- Is this leading order result stable in the perturbation theory?

Natural question

- Is this leading order result stable in the perturbation theory?
- The answer is NO

Natural question

- Is this leading order result stable in the perturbation theory?
- The answer is NO
- Why should we ask this question at all here?

Natural question

- Is this leading order result stable in the perturbation theory?
- The answer is NO
- Why should we ask this question at all here?
- Because we are dealing with partons such as quark and gluons at the initial state which are sensitive to Factorisation scale even at Leading order.

Natural question

- Is this leading order result stable in the perturbation theory?
- The answer is NO
- Why should we ask this question at all here?
- Because we are dealing with partons such as quark and gluons at the initial state which are sensitive to Factorisation scale even at Leading order.

$$d\sigma^{PP}(x, Q^2) = \sum_{ab} \int_x^1 \frac{dz}{z} \Phi_{ab}^{(0)}(z, Q^2, \mu_F^2) \sigma_{ab}^{(0)}\left(\frac{x}{z}, Q^2, M_S^2\right) + \dots$$

Leading Partonic cross section is "independent" of μ_F

Natural question

- Is this leading order result stable in the perturbation theory?
- The answer is NO
- Why should we ask this question at all here?
- Because we are dealing with partons such as quark and gluons at the initial state which are sensitive to Factorisation scale even at Leading order.

$$d\sigma^{PP}(x, Q^2) = \sum_{ab} \int_x^1 \frac{dz}{z} \Phi_{ab}^{(0)}(z, Q^2, \mu_F^2) \sigma_{ab}^{(0)}\left(\frac{x}{z}, Q^2, M_S^2\right) + \dots$$

Leading Partonic cross section is "independent" of μ_F

- Uncertainty can come from Factorisation scale μ_F through the LO flux $\Phi_{ab}^{(0)}(z, \mu_F)$

Natural question

- Is this leading order result stable in the perturbation theory?
- The answer is NO
- Why should we ask this question at all here?
- Because we are dealing with partons such as quark and gluons at the initial state which are sensitive to Factorisation scale even at Leading order.

$$d\sigma^{PP}(x, Q^2) = \sum_{ab} \int_x^1 \frac{dz}{z} \Phi_{ab}^{(0)}(z, Q^2, \mu_F^2) \sigma_{ab}^{(0)}\left(\frac{x}{z}, Q^2, M_S^2\right) + \dots$$

Leading Partonic cross section is "independent" of μ_F

- Uncertainty can come from Factorisation scale μ_F through the LO flux $\Phi_{ab}^{(0)}(z, \mu_F)$
- How serious is it?

Scale Variation of Flux at LHC


$$\Phi_{ab}^I(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a^I(z, \mu_F) f_b^I\left(\frac{x}{z}, \mu_F\right) \quad I = LO, NLO$$

$$\mu_0 = 700 \text{ GeV}, \quad x = \frac{Q}{\sqrt{S}}, \quad Q = 700 \text{ GeV} \quad \sqrt{S} = 14 \text{ TeV}$$

Scale Variation of Flux at LHC

$$\Phi_{ab}^I(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a^I(z, \mu_F) f_b^I\left(\frac{x}{z}, \mu_F\right) \quad I = LO, NLO$$



$$\mu_0 = 700 \text{ GeV}, \quad x = \frac{Q}{\sqrt{S}}, \quad Q = 700 \text{ GeV} \quad \sqrt{S} = 14 \text{ TeV}$$



Scale Variation of Flux at LHC

$$\Phi_{ab}^I(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a^I(z, \mu_F) f_b^I\left(\frac{x}{z}, \mu_F\right) \quad I = LO, NLO$$

$$\mu_0 = 700 \text{ GeV}, \quad x = \frac{Q}{\sqrt{S}}, \quad Q = 700 \text{ GeV} \quad \sqrt{S} = 14 \text{ TeV}$$



Scale Variation of Flux at Tevatron


$$\Phi_{ab}^I(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a^I(z, \mu_F) f_b^I\left(\frac{x}{z}, \mu_F\right) \quad I = LO, NLO$$

$$\mu_0 = 700 \text{ GeV}, \quad x = \frac{Q_0}{\sqrt{S}}, \quad Q = 700 \text{ GeV} \quad (\sqrt{S} = 1.96 \text{ TeV})$$

Scale Variation of Flux at Tevatron

$$\Phi_{ab}^I(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a^I(z, \mu_F) f_b^I\left(\frac{x}{z}, \mu_F\right) \quad I = LO, NLO$$



$$\mu_0 = 700 \text{ GeV}, \quad x = \frac{Q_0}{\sqrt{S}}, \quad Q = 700 \text{ GeV} \quad (\sqrt{S} = 1.96 \text{ TeV})$$



Scale Variation of Flux at Tevatron

$$\Phi_{ab}^I(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a^I(z, \mu_F) f_b^I\left(\frac{x}{z}, \mu_F\right) \quad I = LO, NLO$$

$$\mu_0 = 700 \text{ GeV}, \quad x = \frac{Q_0}{\sqrt{S}}, \quad Q = 700 \text{ GeV} \quad (\sqrt{S} = 1.96 \text{ TeV})$$



We are in Business!

We are in Business!

- Compute Next to leading order **NLO** QCD corrections to **LO** prcesses

We are in Business!

- Compute Next to leading order **NLO** QCD corrections to **LO** processes

$$d\hat{\sigma}_{ab}(\hat{s}, Q^2, \mu_F^2) = d\hat{\sigma}_{ab}^{(0)}(\hat{s}, Q^2, \mu_F^2) \left[1 + \frac{\alpha_s(\mu_R^2)}{4\pi} \Delta_{ab}^{(1)}(\hat{s}, Q^2, \mu_F^2, \mu_R^2) \right]$$

QCD corrections are larger than other EW and gravity corrections.

Standard Model	Gravity
$q + \bar{q} \rightarrow \gamma/Z$ NLO	$q + \bar{q} \rightarrow G$ $g + g \rightarrow G$ NLO

We are in Business!

- Compute Next to leading order **NLO** QCD corrections to **LO** processes

$$d\hat{\sigma}_{ab}(\hat{s}, Q^2, \mu_F^2) = d\hat{\sigma}_{ab}^{(0)}(\hat{s}, Q^2, \mu_F^2) \left[1 + \frac{\alpha_s(\mu_R^2)}{4\pi} \Delta_{ab}^{(1)}(\hat{s}, Q^2, \mu_F^2, \mu_R^2) \right]$$

QCD corrections are larger than other EW and gravity corrections.

Standard Model	Gravity
$q + \bar{q} \rightarrow \gamma/Z$ NLO	$q + \bar{q} \rightarrow G$ $g + g \rightarrow G$ NLO

- Energy momentum tensor is renormalised.

We are in Business!

- Compute Next to leading order **NLO** QCD corrections to **LO** processes

$$d\hat{\sigma}_{ab}(\hat{s}, Q^2, \mu_F^2) = d\hat{\sigma}_{ab}^{(0)}(\hat{s}, Q^2, \mu_F^2) \left[1 + \frac{\alpha_s(\mu_R^2)}{4\pi} \Delta_{ab}^{(1)}(\hat{s}, Q^2, \mu_F^2, \mu_R^2) \right]$$

QCD corrections are larger than other EW and gravity corrections.

Standard Model	Gravity
$q + \bar{q} \rightarrow \gamma/Z$ NLO	$q + \bar{q} \rightarrow G$ $g + g \rightarrow G$ NLO

- Energy momentum tensor is renormalised.
- All the soft and collinear divergences are regulated in dimensional regularisation $n = 4 + \epsilon$.

We are in Business!

- Compute Next to leading order **NLO** QCD corrections to **LO** processes

$$d\hat{\sigma}_{ab}(\hat{s}, Q^2, \mu_F^2) = d\hat{\sigma}_{ab}^{(0)}(\hat{s}, Q^2, \mu_F^2) \left[1 + \frac{\alpha_s(\mu_R^2)}{4\pi} \Delta_{ab}^{(1)}(\hat{s}, Q^2, \mu_F^2, \mu_R^2) \right]$$

QCD corrections are larger than other EW and gravity corrections.


Standard Model	Gravity
$q + \bar{q} \rightarrow \gamma/Z$ NLO	$q + \bar{q} \rightarrow G$ $g + g \rightarrow G$ NLO

- Energy momentum tensor is renormalised.
- All the soft and collinear divergences are regulated in dimensional regularisation $n = 4 + \epsilon$.
- Collinear mass factorisation is done in \overline{MS} scheme.





Virtual Corrections, $q \bar{q} \rightarrow G$

$$\bar{\Delta}_{q\bar{q}}^G = \Delta_{q\bar{q}}^{(0)G} + a_s \frac{2}{\epsilon} \Gamma_{qq}^{(1)} \otimes \Delta_{q\bar{q}}^{(0)G} + a_s \Delta_{q\bar{q}}^{(1)G}$$

$q + \bar{q} \rightarrow G$ (1 loop):







Real emission, $q \bar{q} \rightarrow g G$







Virtual Corrections, $g \bar{g} \rightarrow G$

$$\bar{\Delta}_{gg}^G = \Delta_{gg}^{(0)G} + a_s \frac{2}{\epsilon} \Gamma_{gg}^{(1)} \otimes \Delta_{gg}^{(0)G} + a_s \Delta_{gg}^{(1)G}$$

$g + g \rightarrow G$ (1 loop):







Real emission, $g g \rightarrow g G$



Real emissions, $q \bar{q} \rightarrow q G$

$$\bar{\Delta}_{qg}^G = a_s \frac{1}{\epsilon} \left(\Gamma_{qg}^{(1)} \otimes \Delta_{q\bar{q}}^{(0)G} + \Gamma_{gq}^{(1)} \otimes \Delta_{gg}^{(0)G} \right) + a_s \Delta_{qg}^{(1)G}$$

Real emission, $q g \rightarrow q G$



Invariant mass distribution of lepton pair

Invariant mass distribution of lepton pair

$$\begin{aligned}
 2S \frac{d\sigma^{P_1 P_2}}{dQ^2}(\tau, Q^2) = \\
 \sum_q \mathcal{F}_{SM,q} \left[H_{q\bar{q}}(\tau, Q^2) \otimes \left(\Delta_{q\bar{q}}^{(0)\gamma Z}(\tau, Q^2) + a_s \Delta_{q\bar{q}}^{(1)\gamma Z}(\tau, Q^2) \right) \right. \\
 \left. + \left(H_{qg}(\tau, Q^2) + H_{gq}(\tau, Q^2) \right) \otimes a_s \Delta_{qg}^{(1)\gamma Z}(\tau, Q^2) \right]
 \end{aligned}$$

Invariant mass distribution of lepton pair

$$\begin{aligned}
 2S \frac{d\sigma^{P_1 P_2}}{dQ^2}(\tau, Q^2) = & \\
 & \sum_q \mathcal{F}_{SM,q} \left[H_{q\bar{q}}(\tau, Q^2) \otimes \left(\Delta_{q\bar{q}}^{(0)\gamma Z}(\tau, Q^2) + a_s \Delta_{q\bar{q}}^{(1)\gamma Z}(\tau, Q^2) \right) \right. \\
 & \left. + \left(H_{qg}(\tau, Q^2) + H_{gq}(\tau, Q^2) \right) \otimes a_s \Delta_{qg}^{(1)\gamma Z}(\tau, Q^2) \right] \\
 & + \sum_q \mathcal{F}_G \left[H_{q\bar{q}}(\tau, Q^2) \otimes \left(\Delta_{q\bar{q}}^{(0)G}(\tau, Q^2) + a_s \Delta_{q\bar{q}}^{(1)G}(\tau, Q^2) \right) \right. \\
 & + \left(H_{qg}(\tau, Q^2) + H_{gq}(\tau, Q^2) \right) \otimes a_s \Delta_{qg}^{(1)G}(\tau, Q^2) \\
 & \left. + H_{gg}(\tau, Q^2) \otimes \left(\Delta_{gg}^{(0)G}(\tau, Q^2) + a_s \Delta_{gg}^{(1)G}(\tau, Q^2) \right) \right]
 \end{aligned}$$

Invariant mass distribution of lepton pair

$$\begin{aligned}
 2S \frac{d\sigma^{P_1 P_2}}{dQ^2}(\tau, Q^2) = & \\
 & \sum_q \mathcal{F}_{SM,q} \left[H_{q\bar{q}}(\tau, Q^2) \otimes \left(\Delta_{q\bar{q}}^{(0)\gamma Z}(\tau, Q^2) + a_s \Delta_{q\bar{q}}^{(1)\gamma Z}(\tau, Q^2) \right) \right. \\
 & \left. + \left(H_{qg}(\tau, Q^2) + H_{gq}(\tau, Q^2) \right) \otimes a_s \Delta_{qg}^{(1)\gamma Z}(\tau, Q^2) \right] \\
 & + \sum_q \mathcal{F}_G \left[H_{q\bar{q}}(\tau, Q^2) \otimes \left(\Delta_{q\bar{q}}^{(0)G}(\tau, Q^2) + a_s \Delta_{q\bar{q}}^{(1)G}(\tau, Q^2) \right) \right. \\
 & \left. + \left(H_{qg}(\tau, Q^2) + H_{gq}(\tau, Q^2) \right) \otimes a_s \Delta_{qg}^{(1)G}(\tau, Q^2) \right. \\
 & \left. + H_{gg}(\tau, Q^2) \otimes \left(\Delta_{gg}^{(0)G}(\tau, Q^2) + a_s \Delta_{gg}^{(1)G}(\tau, Q^2) \right) \right]
 \end{aligned}$$

Coefficient Functions Independent of ADD or RS Model

RS Results

	LHC	TEVATRON
\sqrt{S}	14 TeV	1.96 TeV
PDF	MRST 2001	
Choice of scale	LO & NLO $\mu_F = \mu_R$ & $\mu_F = Q$	

RS Results

	LHC	TEVATRON
\sqrt{S}	14 TeV	1.96 TeV
PDF	MRST 2001	
Choice of scale	LO & NLO $\mu_F = \mu_R$ & $\mu_F = Q$	

Distributions:

$$\frac{d\sigma^I(Q)}{dQ}$$

$$\frac{d\sigma^I(Q,Y)}{dQ\,dY}\Big|_{Q_0}$$

RS Results

	LHC	TEVATRON
\sqrt{S}	14 TeV	1.96 TeV
PDF	MRST 2001	
Choice of scale	LO & NLO $\mu_F = \mu_R$ & $\mu_F = Q$	

Distributions:

$$\frac{d\sigma^I(Q)}{dQ} \qquad \frac{d\sigma^I(Q,Y)}{dQ\,dY}\bigg|_{Q_0}$$

K-Factor:

$$K^I \;=\; \left[\frac{d\sigma^I_{LO}(Q)}{dQ}\right]^{-1}\left[\frac{d\sigma^I_{NLO}(Q)}{dQ}\right]$$

RS Results

	LHC	TEVATRON
\sqrt{S}	14 TeV	1.96 TeV
PDF	MRST 2001	
Choice of scale	LO & NLO $\mu_F = \mu_R$ & $\mu_F = Q$	

Distributions:

$$\frac{d\sigma^I(Q)}{dQ} \quad \frac{d\sigma^I(Q, Y)}{dQ dY} \Big|_{Q_0}$$

K-Factor:

$$K^I = \left[\frac{d\sigma_{LO}^I(Q)}{dQ} \right]^{-1} \left[\frac{d\sigma_{NLO}^I(Q)}{dQ} \right]$$

R-Factor:

$$R_{LO}^I = \left[\frac{d\sigma_{LO}^I(Q, \mu = \mu_0)}{dQ} \right]^{-1} \left[\frac{d\sigma_{LO}^I(Q, \mu)}{dQ} \right] \Big|_{Q_0}$$

Partonic Cross Section versus its Flux

- Parton level cross section increases monotonically with invariant lepton pair mass Q upto m_o .

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{RS}(\hat{s}, Q^2, M_S) \sim c_o^4 \lambda \left(\frac{Q^6}{\hat{s} m_o^8} \right), \quad Q < m_o$$

Partonic Cross Section versus its Flux

- Parton level cross section increases monotonically with invariant lepton pair mass Q upto m_o .

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{RS}(\hat{s}, Q^2, M_S) \sim c_o^4 \lambda \left(\frac{Q^6}{\hat{s} m_o^8} \right), \quad Q < m_o$$

- The Parton flux

$$\Phi_{ab} \left(x = \frac{Q^2}{S}, \mu_F \right)$$

decreases as Q or x increases.

Partonic Cross Section versus its Flux

- Parton level cross section increases monotonically with invariant lepton pair mass Q upto m_o .

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{RS}(\hat{s}, Q^2, M_S) \sim c_o^4 \lambda \left(\frac{Q^6}{\hat{s} m_o^8} \right), \quad Q < m_o$$

- The Parton flux

$$\Phi_{ab} \left(x = \frac{Q^2}{S}, \mu_F \right)$$

decreases as Q or x increases.

- At small Q Standard model dominates over Gravity interaction.
- At large Q the gravity "cross section" is comparable to SM.

Partonic Cross Section versus its Flux

- Parton level cross section increases monotonically with invariant lepton pair mass Q upto m_o .

$$\frac{d}{dQ^2} \hat{\sigma}_{ab}^{RS}(\hat{s}, Q^2, M_S) \sim c_o^4 \lambda \left(\frac{Q^6}{\hat{s} m_o^8} \right), \quad Q < m_o$$

- The Parton flux

$$\Phi_{ab} \left(x = \frac{Q^2}{S}, \mu_F \right)$$

decreases as Q or x increases.


- At small Q Standard model dominates over Gravity interaction.
- At large Q the gravity "cross section" is comparable to SM.
- At large Q the parton "cross section" dominates over "Flux" leaving "observable effect".

Flux at LHC and Tevatron

$$\Phi_{ab}(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a(z, \mu_F) f_b\left(\frac{x}{z}, \mu_F\right) \quad x = \frac{Q^2}{S}$$

Flux at LHC and Tevatron


$$\Phi_{ab}(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a(z, \mu_F) f_b\left(\frac{x}{z}, \mu_F\right) \quad x = \frac{Q^2}{S}$$




Gluon flux is largest at LHC

Flux at LHC and Tevatron

$$\Phi_{ab}(x, \mu_F) = \int_x^1 \frac{dz}{z} f_a(z, \mu_F) f_b\left(\frac{x}{z}, \mu_F\right) \quad x = \frac{Q^2}{S}$$



Gluon flux is largest at LHC



Quark-anti quark flux is largest at Tevatron

RS Scenario Results



$$\mathcal{D}(Q^2) = \sum_{n=1}^{\infty} \frac{1}{s - M_n^2 + iM_n\Gamma_n} \equiv \frac{\lambda}{m_0^2}$$

$$\frac{c_0^2}{m_0^2} \mathcal{D}(Q^2) = \frac{c_0^2}{m_0^4} \lambda$$

RS Scenario Results


$$\mathcal{D}(Q^2) = \sum_{n=1}^{\infty} \frac{1}{s - M_n^2 + iM_n\Gamma_n} \equiv \frac{\lambda}{m_0^2}$$

$$\frac{c_0^2}{m_0^2} \mathcal{D}(Q^2) = \frac{c_0^2}{m_0^4} \lambda$$



- Away from the resonance region gravity contribution is negligible
- K-Factor behavior can be understood from the $K^{(0)}$ behavior for the RS model.

Rapidity Y distribution:




- K-Factor for rapidity distribution, close to to the first KK resonance $m_1 \sim 1.5$ TeV
- Gravity dominates the resonance region as can be understood from the $K^{(0)}$ behavior of RS model.

K-Factor

$$K^{(SM+GR)}(Q) = \frac{K^{SM} + K^{GR} G^{(0)}}{1 + G^{(0)}}$$



$$G^{(0)}(Q) = \left[\frac{d\sigma_{LO}^{SM}(Q)}{dQ} \right]^{-1} \left[\frac{d\sigma_{LO}^{GR}(Q)}{dQ} \right]$$



- $G^{(0)}(Q)$ behavior is governed by a competing 'couplings' and PDF flux at LHC and TEV
- At high Q when Gravity contribution becomes comparable to SM, the PDF flux dictates the proceedings

R-Factor:



$$R_{LO,NLO}^I = \left[\frac{d\sigma_{LO,NLO}^I(Q, \mu = \mu_0)}{dQ} \right]^{-1} \left[\frac{d\sigma_{LO,NLO}^I(Q, \mu)}{dQ} \right] \Big|_{Q=Q_0}$$



- Scale variation reduced considerably in going from LO \rightarrow NLO
- Inclusion of SM to GR also reduces scale variation

RS Scenario Results



$$\frac{c_0^2}{m_0^2} \mathcal{D}(Q^2) = \frac{c_0^2}{m_0^2} \sum_{n=1}^{\infty} \frac{1}{s - M_n^2 + iM_n\Gamma_n} \equiv \frac{c_0^2}{m_0^4} \lambda\left(\frac{Q}{m_0}\right)$$



- Away from the resonance region gravity contribution is negligible
- K-Factor behavior can be understood from the $K^{(0)}$ behavior for the RS model

Angular distribution:



$$\left. \frac{d\sigma(Q, \cos \theta)}{dQ d\cos \theta} \right|_{Q_0}$$





Factorisation scale dependence of angular distribution:

$$\left. \frac{d\sigma(Q, \cos\theta)}{dQ d\cos\theta} \right|_{Q_0}$$

Distributions	Tevatron		LHC	
	LO	NLO	LO	NLO
$d^2\sigma/dQdY$	23.2	7.7	18.7	6.9
$d^2\sigma/dQd\cos\theta$	24.2	8.0	18.4	6.8



RS Rapidity



Summary

- Next to Leading Order coefficient functions for DY process in models of TeV-scale gravity are available now.
- Various distributions *viz.* Q , x_F , Y distributions and A_{FB} asymmetry at NLO are studied for ADD & RS models.
- Theoretical uncertainties get significantly reduced at NLO level
- Quantitative impact of the QCD corrections for searches of extra dimension at hadron colliders investigated