

Phenomenology of two Universal Extra Dimensions

Kirtiman Ghosh

Calcutta University, Kolkata, India

Based on:

KG and A. Datta [arXiv:0801.0943 and 0802.2162(hep-ph)]

Plan of talk

- Introduction to *2-Universal Extra Dimensions (2-UED)*
- *Particle Spectra*
- *Spinless Adjoints* and *Interactions*
- Production of *Spinless Adjoints* at the LHC.
- Signature of EW *KK Gauge Bosons* at the ILC.
- Concluding Remarks

Introduction to *2-UED*

As the name suggests, in *2UED* all the *SM fields* can propagate universally in the *six-dimensional* space-time x^α , $\alpha = 0, \dots, 5$.

- x^μ ($\mu = 0, 1, 2, 3$) form the usual *Minkowski space*.

Introduction to *2-UED*


As the name suggests, in *2UED* all the *SM fields* can propagate universally in the *six-dimensional* space-time x^α , $\alpha = 0, \dots, 5$.

- x^μ ($\mu = 0, 1, 2, 3$) form the usual *Minkowski space*.
- x^4 and x^5 are transverse and flat, compactified with $0 \leq x^4, x^5 \leq L$.

Introduction to *2-UED*


As the name suggests, in *2UED* all the *SM fields* can propagate universally in the *six-dimensional* space-time x^α , $\alpha = 0, \dots, 5$.

- x^μ ($\mu = 0, 1, 2, 3$) form the usual *Minkowski space*.
- x^4 and x^5 are transverse and flat, compactified with $0 \leq x^4, x^5 \leq L$.




Compactification

- Toroidal Compactification




Compactification

- Toroidal Compactification
 - leads to vector-like 4D fermions




Compactification

- Toroidal Compactification
 - leads to vector-like 4D fermions
- Next alternative: fold the square along a diagonal.
 $(y, 0) \equiv (0, y), \quad (y, L) \equiv (L, y), \quad \forall y \in [0, L]$



Compactification

- **Toroidal Compactification**
 - leads to **vector-like** 4D fermions
- Next alternative: **fold the square along a diagonal.**
 - Leaves at least a single 4D fermion of definite chirality.



Continued...

The physics at **identified points** is identical if the *Lagrangian* takes the same value for any field configuration:

$$\mathcal{L}|_{x^\mu, y, 0} = \mathcal{L}|_{x^\mu, 0, y}; \quad \mathcal{L}|_{x^\mu, y, L} = \mathcal{L}|_{x^\mu, L, y}$$

- This requirement fixes the **boundary conditions (BC)** for 6D scalar fields $\Phi(x^\alpha)$ and 6D Weyl fermions $\Psi_\pm(x^\alpha)$.

Continued...

The physics at **identified points** is identical if the *Lagrangian* takes the same value for any field configuration:

$$\mathcal{L}|_{x^\mu, y, 0} = \mathcal{L}|_{x^\mu, 0, y}; \quad \mathcal{L}|_{x^\mu, y, L} = \mathcal{L}|_{x^\mu, L, y}$$

- This requirement fixes the **boundary conditions (BC)** for 6D scalar fields $\Phi(x^\alpha)$ and 6D Weyl fermions $\Psi_\pm(x^\alpha)$.
- **Gauge Symmetry** + BC for scalar fields $\Phi(x^\alpha)$ or Weyl fermions $\Psi_\pm(x^\alpha)$.

Continued...

The physics at **identified points** is identical if the *Lagrangian* takes the same value for any field configuration:

$$\mathcal{L}|_{x^\mu, y, 0} = \mathcal{L}|_{x^\mu, 0, y}; \quad \mathcal{L}|_{x^\mu, y, L} = \mathcal{L}|_{x^\mu, L, y}$$

- This requirement fixes the **boundary conditions (BC)** for 6D scalar fields $\Phi(x^\alpha)$ and 6D Weyl fermions $\Psi_\pm(x^\alpha)$.
- **Gauge Symmetry** + BC for scalar fields $\Phi(x^\alpha)$ or Weyl fermions $\Psi_\pm(x^\alpha)$.
 - Fixes the BC's for 6D gauge fields.

Decomposition

Any 6D field (fermion/gauge or scalar) $\Phi(x^\mu, x^4, x^5)$ can be decomposed as:

$$\Phi(x^\mu, x^4, x^5) = \frac{1}{L} \sum_{j,k} f_n^{(j,k)}(x^4, x^5) \Phi^{(j,k)}(x^\mu)$$

$$f_n^{(j,k)}(x^4, x^5) = \frac{1}{1 + \delta_{j,0}\delta_{k,0}} \left[e^{-in\pi/2} \cos\left(\frac{jx^4 + kx^5}{R} + \frac{n\pi}{2}\right) + \cos\left(\frac{kx^4 - jx^5}{R} + \frac{n\pi}{2}\right) \right]$$

- $\Phi^{(j,k)}(x^\mu)$ are the KK modes of 6D field $\Phi(x^\mu, x^4, x^5)$ with KK-parity $(-1)^{j+k}$

Decomposition

Any 6D field (fermion/gauge or scalar) $\Phi(x^\mu, x^4, x^5)$ can be decomposed as:

$$\Phi(x^\mu, x^4, x^5) = \frac{1}{L} \sum_{j,k} f_n^{(j,k)}(x^4, x^5) \Phi^{(j,k)}(x^\mu)$$

$$f_n^{(j,k)}(x^4, x^5) = \frac{1}{1 + \delta_{j,0}\delta_{k,0}} \left[e^{-in\pi/2} \cos\left(\frac{jx^4 + kx^5}{R} + \frac{n\pi}{2}\right) + \cos\left(\frac{kx^4 - jx^5}{R} + \frac{n\pi}{2}\right) \right]$$

- $\Phi^{(j,k)}(x^\mu)$ are the KK modes of 6D field $\Phi(x^\mu, x^4, x^5)$ with KK-parity $(-1)^{j+k}$
- n is an integer whose value is restricted to 0, 1, 2 or 3 by the BC. And $j + l/2$ and $k + l/2$ are integers with $l = 0$ or 1.

Decomposition

Any 6D field (fermion/gauge or scalar) $\Phi(x^\mu, x^4, x^5)$ can be decomposed as:

$$\Phi(x^\mu, x^4, x^5) = \frac{1}{L} \sum_{j,k} f_n^{(j,k)}(x^4, x^5) \Phi^{(j,k)}(x^\mu)$$

$$f_n^{(j,k)}(x^4, x^5) = \frac{1}{1 + \delta_{j,0}\delta_{k,0}} \left[e^{-in\pi/2} \cos\left(\frac{jx^4 + kx^5}{R} + \frac{n\pi}{2}\right) + \cos\left(\frac{kx^4 - jx^5}{R} + \frac{n\pi}{2}\right) \right]$$

- $\Phi^{(j,k)}(x^\mu)$ are the KK modes of 6D field $\Phi(x^\mu, x^4, x^5)$ with KK-parity $(-1)^{j+k}$
- n is a integer whose value is restricted to 0, 1, 2 or 3 by the BC. And $j + l/2$ and $k + l/2$ are integers with $l = 0$ or 1.
 - There are only eight self-consistent choices of BC.

Decomposition

Any 6D field (fermion/gauge or scalar) $\Phi(x^\mu, x^4, x^5)$ can be decomposed as:

$$\Phi(x^\mu, x^4, x^5) = \frac{1}{L} \sum_{j,k} f_n^{(j,k)}(x^4, x^5) \Phi^{(j,k)}(x^\mu)$$

$$f_n^{(j,k)}(x^4, x^5) = \frac{1}{1 + \delta_{j,0}\delta_{k,0}} \left[e^{-in\pi/2} \cos\left(\frac{jx^4 + kx^5}{R} + \frac{n\pi}{2}\right) + \cos\left(\frac{kx^4 - jx^5}{R} + \frac{n\pi}{2}\right) \right]$$

- $\Phi^{(j,k)}(x^\mu)$ are the KK modes of 6D field $\Phi(x^\mu, x^4, x^5)$ with KK-parity $(-1)^{j+k}$
- n is a **integer** whose value is restricted to 0, 1, 2 or 3 by the BC. And $j + l/2$ and $k + l/2$ are **integers** with $l = 0$ or 1.
 - There are only **eight** self-consistent choices of BC.
- Form of $f_n^{(j,k)}(x^4, x^5)$ tells that only $n = 0$ and $l = 0$ allows zero mode ($j = k = 0$) fields in the **4D effective theory**.

Fields in 6D

- **Spin-1/2 fields** in 6D:

- There six anti-commuting matrices Γ^α , $\alpha = 1, \dots, 5$ with minimum dimensionality 8×8 , defined as $\Gamma^\mu = \gamma^\mu \otimes \sigma^0$, $\Gamma^{4,5} = i\gamma_5 \otimes \sigma^{1,2}$
- 6D fermions have $+$ or $-$ chirality, defined by the eigenvalue of the $\bar{\Gamma} = -\gamma_5 \otimes \sigma^3$ 6D chirality operator: $\bar{\Gamma}\Psi_\pm = \pm \Psi_\pm$.
- A 6D chiral fermion include both 4D chiralities: $\gamma_5 \otimes \sigma^0 \Psi_{\pm L} = -\Psi_{\pm L}$ and $\gamma_5 \otimes \sigma^0 \Psi_{\pm R} = -\Psi_{\pm R}$
- BC for Left and Right chiral components 6D Weyl fermions are related by:
 $n_L^\pm - n_R^\pm = \pm 1$

- **Spin-1 fields** in 6D:

- In 6D, the gauge fields has six components $A_\alpha(x^\mu, x^4, x^5)$, $\alpha = 0, \dots, 5$.
- Upon compactification, they decompose into towers of 4D spin-1 fields $A_\nu^{(j,k)}(x^\mu)$, two towers of spin-0 fields $A_4^{(j,k)}(x^\mu)$ and $A_5^{(j,k)}(x^\mu)$.

Continued...

- $A_{4,5}^{(j,k)}(x^\mu)$ are not the physical fields. Define at each KK label two scalar fields, $A_G^{(j,k)}$ and $A_H^{(j,k)}$ such that:

$$\partial_4 A_5 - \partial_5 A_4 = \frac{1}{L} \sum_{j \geq 1} \sum_{k \geq 0} M_{(j,k)} A_H^{(j,k)}(x^\nu) f_0^{(j,k)}(x^4, x^5)$$

$$\partial_4 A_4 + \partial_5 A_5 = \frac{1}{L} \sum_{j \geq 1} \sum_{k \geq 0} M_{(j,k)} A_G^{(j,k)}(x^\nu) f_0^{(j,k)}(x^4, x^5)$$

- Under 6D gauge transformations: $A_\alpha \rightarrow A_\alpha + \partial_\alpha \xi / g_6$, $A_H^{(j,k)}$ remains invariant but $A_G^{(j,k)}$ does not.

• Spin-0 fields in 6D

- $\mathcal{L}_\Phi = |D_\alpha \Phi|^2 - \frac{\lambda_6}{2} (\Phi^\dagger \Phi - \frac{1}{2} v_6^2)^2$ with $\Phi = \frac{1}{\sqrt{2}} (v_6 + h) e^{i \frac{\eta}{v_6}}$
 - Compactification: $h^{(j,k)}(x^\mu)$ and $\eta^{(j,k)}(x^\mu)$, (j, k) includes $(0, 0)$.
 - Mixing with **Spinless Adjoints** And **Physical fields**:
 - $h^{(j,k)}(x^\mu)$: KK tower of SM Higgs, $M_h^{(j,k)} = \sqrt{M_{j,k}^2 + \lambda_4 v_4^2}$
 - $A_H^{(j,k)}(x^\mu)$: KK tower of **Spinless Adjoints**, $M_{A_H}^{(j,k)} = \sqrt{M_{j,k}^2 + g_4^2 v_4^2}$
- $$\tilde{A}_G^{(j,k)} = \left(M_{j,k} A_G^{(j,k)} + g_4 v_6 \eta^{(j,k)} \right) / M_A^{(j,k)}$$

The SM in 6D

In 6D, the fields and BC's of the fields are chosen such that upon compactification the **Zero-Modes** reproduce SM.

- Anomaly cancellation + fermion mass generation:
Force the choice of quarks generations to be
 $Q_+ \equiv (U_+, D_+), U_-, D_-$.

The SM in 6D

In 6D, the fields and BC's of the fields are chosen such that upon compactification the **Zero-Modes** reproduce SM.

- Anomaly cancellation + fermion mass generation:
Force the choice of quarks generations to be
 $Q_+ \equiv (U_+, D_+), U_-, D_-$.
- After **compactification**

The SM in 6D

In 6D, the fields and BC's of the fields are chosen such that upon compactification the **Zero-Modes** reproduce SM.

- Anomaly cancellation + fermion mass generation:
Force the choice of quarks generations to be

$$Q_+ \equiv (U_+, D_+), U_-, D_-.$$

- After **compactification**

- 6D doublet fermions $\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_+$: $\begin{pmatrix} \psi_1^{(j,k)} \\ \psi_2^{(j,k)} \end{pmatrix}_{+L} + \begin{pmatrix} \psi_1^{(j,k)} \\ \psi_2^{(j,k)} \end{pmatrix}_{+R} + \begin{pmatrix} \psi_1^{(0,0)} \\ \psi_2^{(0,0)} \end{pmatrix}_{+L}$

The SM in 6D

In 6D, the fields and BC's of the fields are chosen such that upon compactification the **Zero-Modes** reproduce SM.

- Anomaly cancellation + fermion mass generation:
Force the choice of quarks generations to be

$$Q_+ \equiv (U_+, D_+), U_-, D_-.$$

- After **compactification**

- 6D doublet fermions $\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_+$: $\begin{pmatrix} \psi_1^{(j,k)} \\ \psi_2^{(j,k)} \end{pmatrix}_{+L} + \begin{pmatrix} \psi_1^{(j,k)} \\ \psi_2^{(j,k)} \end{pmatrix}_{+R} + \begin{pmatrix} \psi_1^{(0,0)} \\ \psi_2^{(0,0)} \end{pmatrix}_{+L}$
- 6D singlet fermions ψ_- : $\psi_{-L}^{(j,k)} + \psi_{-R}^{(j,k)} + \psi_{-R}^{(0,0)}$

The SM in 6D

In 6D, the fields and BC's of the fields are chosen such that upon compactification the **Zero-Modes** reproduce SM.

- Anomaly cancellation + fermion mass generation:
Force the choice of quarks generations to be

$$Q_+ \equiv (U_+, D_+), U_-, D_-.$$

- After **compactification**

- 6D doublet fermions $\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_+$: $\begin{pmatrix} \psi_1^{(j,k)} \\ \psi_2^{(j,k)} \end{pmatrix}_{+L} + \begin{pmatrix} \psi_1^{(j,k)} \\ \psi_2^{(j,k)} \end{pmatrix}_{+R} + \begin{pmatrix} \psi_1^{(0,0)} \\ \psi_2^{(0,0)} \end{pmatrix}_{+L}$
- 6D singlet fermions ψ_- : $\psi_{-L}^{(j,k)} + \psi_{-R}^{(j,k)} + \psi_{-R}^{(0,0)}$
- These **zero mode** fields are identified with the **SM fermions**. As for example,
 $(u_L, d_L) \equiv Q_{+L}^{(0,0)}(x^\mu)$, $u_R \equiv U_{-R}^{(0,0)}(x^\mu)$ and $d_R \equiv D_{-R}^{(0,0)}(x^\mu)$.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
 - 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
 - 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.
 - 6D $U(1)$ gauge boson $B_\alpha(x^\beta)$: $B_\mu^{(j,k)}(x^\nu)$ and $B_H^{(j,k)}(x^\nu)$.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.

- 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
- 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.
- 6D $U(1)$ gauge boson $B_\alpha(x^\beta)$: $B_\mu^{(j,k)}(x^\nu)$ and $B_H^{(j,k)}(x^\nu)$.
- $G_\mu^{a(0,0)}$ and $W_\mu^{\pm(0,0)}$: Identified with SM gluons and W -bosons.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
 - 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.
 - 6D $U(1)$ gauge boson $B_\alpha(x^\beta)$: $B_\mu^{(j,k)}(x^\nu)$ and $B_H^{(j,k)}(x^\nu)$.
 - $G_\mu^{a(0,0)}$ and $W_\mu^{\pm(0,0)}$: Identified with SM gluons and W-bosons.
 - SM mixing of $W_\mu^{3(0,0)}$ and $B_\mu^{(0,0)}$: Gives photon and Z-boson.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
 - 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.
 - 6D $U(1)$ gauge boson $B_\alpha(x^\beta)$: $B_\mu^{(j,k)}(x^\nu)$ and $B_H^{(j,k)}(x^\nu)$.
 - $G_\mu^{a(0,0)}$ and $W_\mu^{\pm(0,0)}$: Identified with SM gluons and W-bosons.
 - SM mixing of $W_\mu^{3(0,0)}$ and $B_\mu^{(0,0)}$: Gives photon and Z-boson.
 - For non-zero modes, mixing of $W_\mu^{3(j,k)}$, $B_\mu^{(j,k)}$ as well as $W_H^{3(j,k)}$, $B_H^{(j,k)}$ is small.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
 - 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.
 - 6D $U(1)$ gauge boson $B_\alpha(x^\beta)$: $B_\mu^{(j,k)}(x^\nu)$ and $B_H^{(j,k)}(x^\nu)$.
 - $G_\mu^{a(0,0)}$ and $W_\mu^{\pm(0,0)}$: Identified with SM gluons and W-bosons.
 - SM mixing of $W_\mu^{3(0,0)}$ and $B_\mu^{(0,0)}$: Gives photon and Z-boson.
 - For non-zero modes, mixing of $W_\mu^{3(j,k)}$, $B_\mu^{(j,k)}$ as well as $W_H^{3(j,k)}$, $B_H^{(j,k)}$ is small.
- **6D Higgs Doublet:** A tower of 4D Scalar Weak Doublets.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
 - 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.
 - 6D $U(1)$ gauge boson $B_\alpha(x^\beta)$: $B_\mu^{(j,k)}(x^\nu)$ and $B_H^{(j,k)}(x^\nu)$.
 - $G_\mu^{a(0,0)}$ and $W_\mu^{\pm(0,0)}$: Identified with SM gluons and W-bosons.
 - SM mixing of $W_\mu^{3(0,0)}$ and $B_\mu^{(0,0)}$: Gives photon and Z-boson.
 - For non-zero modes, mixing of $W_\mu^{3(j,k)}$, $B_\mu^{(j,k)}$ as well as $W_H^{3(j,k)}$, $B_H^{(j,k)}$ is small.
- **6D Higgs Doublet:** A tower of 4D Scalar Weak Doublets.
 - Zero-mode doublet: W and Z-boson mass + and the SM Higgs.

continued...

- **6D Gauge Bosons:** Tower of 4D Gauge bosons + 4D Scalars.
 - 6D $SU(3)$ gauge bosons $G_\alpha^a(x^\beta)$: $G_\mu^{a(j,k)}(x^\nu)$ and $G_H^{a(j,k)}(x^\nu)$.
 - 6D $SU(2)$ gauge bosons $W_\alpha^a(x^\beta)$: $W_\mu^{3(j,k)}(x^\nu)$ and $W_H^{3(j,k)}(x^\nu)$, $W_\mu^{\pm(j,k)}(x^\nu)$, $W_H^{\pm(j,k)}(x^\nu)$.
 - 6D $U(1)$ gauge boson $B_\alpha(x^\beta)$: $B_\mu^{(j,k)}(x^\nu)$ and $B_H^{(j,k)}(x^\nu)$.
 - $G_\mu^{a(0,0)}$ and $W_\mu^{\pm(0,0)}$: Identified with SM gluons and W-bosons.
 - SM mixing of $W_\mu^{3(0,0)}$ and $B_\mu^{(0,0)}$: Gives photon and Z-boson.
 - For non-zero modes, mixing of $W_\mu^{3(j,k)}$, $B_\mu^{(j,k)}$ as well as $W_H^{3(j,k)}$, $B_H^{(j,k)}$ is small.
- **6D Higgs Doublet:** A tower of 4D Scalar Weak Doublets.
 - Zero-mode doublet: W and Z-boson mass + and the SM Higgs.
 - KK-doublet: KK tower for SM Higgs + mix with longitudinal modes of KK EW gauge bosons. (mixing is suppressed by R^{-1}).

6UED *Lagrangian*

- Complete 4D effective *Lagrangian*:

$$\begin{aligned}\mathcal{L}_{4D} = & \int_0^L dx^4 \int_0^L dx^5 \mathcal{L}_{bulk} + \delta(x_4)\delta(L-x_5)\mathcal{L}_2 + [\delta(x_4)\delta(x_5) + \\ & + \delta(L-x_4)\delta(L-x_5)]\mathcal{L}_1\end{aligned}$$

6UED *Lagrangian*

- Complete 4D effective *Lagrangian*:

$$\begin{aligned}\mathcal{L}_{4D} = & \int_0^L dx^4 \int_0^L dx^5 \mathcal{L}_{bulk} + \delta(x_4)\delta(L-x_5)\mathcal{L}_2 + [\delta(x_4)\delta(x_5) + \\ & + \delta(L-x_4)\delta(L-x_5)]\mathcal{L}_1\end{aligned}$$

- \mathcal{L}_{bulk} includes SM like interactions in 6D.

6UED *Lagrangian*

- Complete 4D effective *Lagrangian*:

$$\begin{aligned}\mathcal{L}_{4D} = & \int_0^L dx^4 \int_0^L dx^5 \mathcal{L}_{bulk} + \delta(x_4)\delta(L - x_5)\mathcal{L}_2 + [\delta(x_4)\delta(x_5) + \\ & + \delta(L - x_4)\delta(L - x_5)]\mathcal{L}_1\end{aligned}$$

- \mathcal{L}_{bulk} includes SM like interactions in 6D.
- \mathcal{L}_1 and \mathcal{L}_2 contain *Localized Operators*

6UED *Lagrangian*

- Complete 4D effective *Lagrangian*:

$$\begin{aligned}\mathcal{L}_{4D} = & \int_0^L dx^4 \int_0^L dx^5 \mathcal{L}_{bulk} + \delta(x_4)\delta(L - x_5)\mathcal{L}_2 + [\delta(x_4)\delta(x_5) + \\ & + \delta(L - x_4)\delta(L - x_5)]\mathcal{L}_1\end{aligned}$$

- \mathcal{L}_{bulk} includes SM like interactions in 6D.
- \mathcal{L}_1 and \mathcal{L}_2 contain *Localized Operators*
- Contributions to those *Localized Operators*:

6UED *Lagrangian*

- Complete 4D effective *Lagrangian*:

$$\begin{aligned}\mathcal{L}_{4D} = & \int_0^L dx^4 \int_0^L dx^5 \mathcal{L}_{bulk} + \delta(x_4)\delta(L-x_5)\mathcal{L}_2 + [\delta(x_4)\delta(x_5) + \\ & + \delta(L-x_4)\delta(L-x_5)]\mathcal{L}_1\end{aligned}$$

- \mathcal{L}_{bulk} includes SM like interactions in 6D.
- \mathcal{L}_1 and \mathcal{L}_2 contain *Localized Operators*
- Contributions to those *Localized Operators*:
 - **Loop Effect.**

6UED *Lagrangian*

- Complete 4D effective *Lagrangian*:

$$\begin{aligned}\mathcal{L}_{4D} = & \int_0^L dx^4 \int_0^L dx^5 \mathcal{L}_{bulk} + \delta(x_4)\delta(L - x_5)\mathcal{L}_2 + [\delta(x_4)\delta(x_5) + \\ & + \delta(L - x_4)\delta(L - x_5)]\mathcal{L}_1\end{aligned}$$

- \mathcal{L}_{bulk} includes SM like interactions in 6D.
- \mathcal{L}_1 and \mathcal{L}_2 contain *Localized Operators*
- Contributions to those *Localized Operators*:
 - Loop Effect.
 - Physics above the cut-off scale.

Mass Spectrum

The **tree-level masses** for (j, k) -th KK-mode particles are given by $\sqrt{M_{j,k}^2 + m_0^2}$, where $M_{j,k} = \sqrt{j^2 + k^2}/R$. m_0 is the mass of the corresponding zero mode particle.

- Tree-level masses are approximately **degenerate**.

Mass Spectrum


The **tree-level masses** for (j, k) -th KK-mode particles are given by $\sqrt{M_{j,k}^2 + m_0^2}$, where $M_{j,k} = \sqrt{j^2 + k^2}/R$. m_0 is the mass of the corresponding zero mode particle.

- Tree-level masses are approximately **degenerate**.
- **Degeneracy** is lifted by **radiative effects**.

Mass Spectrum

The **tree-level masses** for (j, k) -th KK-mode particles are given by $\sqrt{M_{j,k}^2 + m_0^2}$, where $M_{j,k} = \sqrt{j^2 + k^2}/R$. m_0 is the mass of the corresponding zero mode particle.

- Tree-level masses are approximately **degenerate**.
- **Degeneracy** is lifted by **radiative effects**.







G. Burdman, B. Dobrescu, E. Ponton, [Phys. Rev. D **74**, 075008 (2006)]

Interactions of Spinless Adjoints

- KK-number conserving (KKNC) interactions.

$$\left[i \int_0^L dx^4 \int_0^L dx^5 (\bar{\Psi}_{\pm} \Gamma^{\alpha} D_{\alpha} \Psi_{\pm}) \right]$$



Interactions (KKNV)

Localized Operators, after compactification, give rise to the KK number violating (KKNV) 2-point and 3-point functions.

- Introduce kinetic and mass mixing between KK states.

Interactions (KKNV)

Localized Operators, after compactification, give rise to the KK number violating (KKNV) 2-point and 3-point functions.

- Introduce kinetic and mass mixing between KK states.
- Mixing Effect + KKNV 3-point function: Induce KKNV couplings.

Interactions (KKNV)

Localized Operators, after compactification, give rise to the KK number violating (KKNV) 2-point and 3-point functions.

- Introduce kinetic and mass mixing between KK states.
- Mixing Effect + KKNV 3-point function: Induce KKNV couplings.
- For Example: Coupling involving two zero mode fermions and

Interactions (KKNV)

Localized Operators, after compactification, give rise to the KK number violating (KKNV) 2-point and 3-point functions.

- Introduce kinetic and mass mixing between KK states.
- Mixing Effect + KKNV 3-point function: Induce KKNV couplings.
- For Example: Coupling involving two zero mode fermions and

- KK gauge boson $C_{j,k}^{f_R A} \left[\bar{\psi}_{-R}^{(0,0)} \gamma^\mu A_\mu^{(j,k)} \psi_{-R}^{(0,0)} \right]$

Interactions (KKNV)

Localized Operators, after compactification, give rise to the KK number violating (KKNV) 2-point and 3-point functions.



- Introduce kinetic and mass mixing between KK states.
- Mixing Effect + KKNV 3-point function: Induce KKNV couplings.
- For Example: Coupling involving two zero mode fermions and

- KK gauge boson $C_{j,k}^{f_R A} \left[\bar{\psi}_{-R}^{(0,0)} \gamma^\mu A_\mu^{(j,k)} \psi_{-R}^{(0,0)} \right]$
- Spinless Adjoint $\frac{\tilde{C}_{j,k}^{f_R A}}{M_{(j,k)}} \left[\bar{\psi}_{-R}^{(0,0)} \gamma^\mu \psi_{-R}^{(0,0)} \right] \partial_\mu A_H^{(j,k)}$

KG, A. Datta [arXiv:0801.0943(hep-ph)]

Interactions (KKNV)

- There are another class of operator arise from finite 1-loop effect.





Interactions (KKNV)

- There are another class of operator arise from finite 1-loop effect.

- Resulting **Dimension-5 operator**:

$$C_{AB}^V \epsilon^{\mu\nu\alpha\beta} \left[A_{\mu\nu}^{(J_2, K_2)} B_{\alpha\beta} V_H^{(J_1, K_1)} \right]$$





Interactions (KKNV)

- There are another class of operator arise from finite 1-loop effect.

- Resulting **Dimension-5 operator**:

$$C_{AB}^V \epsilon^{\mu\nu\alpha\beta} \left[A_{\mu\nu}^{(J_2, K_2)} B_{\alpha\beta} V_H^{(J_1, K_1)} \right]$$

- **Anomaly Cancellation.**





Interactions (KKNV)

- There are another class of operator arise from finite 1-loop effect.

- Resulting **Dimension-5 operator**:

$$C_{AB}^V \epsilon^{\mu\nu\alpha\beta} \left[A_{\mu\nu}^{(J_2, K_2)} B_{\alpha\beta} V_H^{(J_1, K_1)} \right]$$

- **Anomaly Cancellation.**
- Mass Splitting between KK fermions plays very crucial rule.




Interactions (KKNV)

- There are another class of operator arise from finite 1-loop effect.

- Resulting **Dimension-5 operator**:

$$C_{AB}^V \epsilon^{\mu\nu\alpha\beta} \left[A_{\mu\nu}^{(J_2, K_2)} B_{\alpha\beta} V_H^{(J_1, K_1)} \right]$$




- Anomaly Cancellation.**




- Mass Splitting between KK fermions plays very crucial rule.

- Dimension-6 operator**:

$$\left[B_{\mu\nu} \partial^\mu A_H^{(J_1, K_1)} \partial^\nu V_H^{(J_2, K_2)} \right]$$



Decays of (1,1) Spinless Adjoints




Single production at LHC

- Coupling with light **SM quark** is suppressed by **quark mass**.

Single production at LHC



- Coupling with light **SM quark** is suppressed by **quark mass**.
- Coupling with **SM gluons** generated by finite **1-Loop Effect**.

$$\sigma(pp \rightarrow V_H^{(1,1)} + X) = \frac{\pi^2}{36sm_{V_H}} \Gamma(V_H^{(1,1)} \rightarrow gg) \int_{\tau}^1 \frac{dx}{x} g(x, m_V^2) g\left(\frac{\tau}{x}, m_V^2\right)$$



Pair production at LHC



Pair production of **Spinless Adjoints** takes place via **KKNC**.



- Being a **pure QCD** process $G_H^{(1,1)}$ has a large cross-section at the LHC.

Pair production at LHC

Pair production of **Spinless Adjoints** takes place via **KKNC**.



- Being a **pure QCD** process $G_H^{(1,1)}$ has a large cross-section at the LHC.
- Pair productions of **EW Spinless Adjoint** are miniscule even for lower values of R^{-1} .

Signatures at future e^+e^- collider



Even KK parity **gauge bosons** couples with e^+e^- via **KKNV** interactions.

- We consider $B_\mu^{(1,1)}$, $W_\mu^{3(1,1)}$ production in association with a **photon**. This has some interesting consequences.

Signatures at future e^+e^- collider

Even KK parity **gauge bosons** couples with e^+e^- via **KKNV** interactions.

- We consider $B_\mu^{(1,1)}$, $W_\mu^{3(1,1)}$ production in association with a **photon**. This has some interesting consequences.
 - **Production cross-section** grows with mass of $B_\mu^{(1,1)}$ or $W_\mu^{3(1,1)}$.



KG, A. Datta [arXiv:0802.2162(hep-ph)]

Continued...

- **Photon** energy $E_\gamma = \frac{s - m_{V_\mu}^2}{2\sqrt{s}}$.

Continued...

- **Photon** energy $E_\gamma = \frac{s - m_{V_\mu}^2}{2\sqrt{s}}$.
 - **Photon** energy distribution is ideally an **delta-function**.

Continued...



- **Photon** energy $E_\gamma = \frac{s - m_{V_\mu}^2}{2\sqrt{s}}$.
 - **Photon** energy distribution is ideally an **delta-function**.
 - Due to finite detector resolution and **initial state radiations (ISR)** the **photon** energy distribution is **smeared**.

Continued...



- **Photon** energy $E_\gamma = \frac{s - m_{V_\mu}^2}{2\sqrt{s}}$.
 - **Photon** energy distribution is ideally an **delta-function**.
 - Due to finite detector resolution and **initial state radiations (ISR)** the **photon** energy distribution is **smeared**.
 - But, the **position** of the peak does not shift.

Continued...

- **Photon** energy $E_\gamma = \frac{s - m_{V_\mu}^2}{2\sqrt{s}}$.
 - **Photon** energy distribution is ideally an **delta-function**.
 - Due to finite detector resolution and **initial state radiations (ISR)** the **photon** energy distribution is **smeared**.
 - But, the **position** of the peak does not shift.
 - $B_\mu^{(1,1)}$ or $W_\mu^{3(1,1)}$ mass: Measurable from mono-energetic **photon**.





Continued...





- **Signals** are always well above the 5σ fluctuations of the **backgrounds**. This opens up a possibility, to measure the relative strengths of the signals in $2j$ and $2t$ channel.

Continued...





- **Signals** are always well above the 5σ fluctuations of the **backgrounds**. This opens up a possibility, to measure the relative strengths of the signals in $2j$ and $2t$ channel.
 - One can determine $\frac{\Gamma(B_\mu^{(1,1)} \rightarrow 2j)}{\Gamma(B_\mu^{(1,1)} \rightarrow 2t)}$

Continued...





- **Signals** are always well above the 5σ fluctuations of the **backgrounds**. This opens up a possibility, to measure the relative strengths of the signals in $2j$ and $2t$ channel.
 - One can determine $\frac{\Gamma(B_{\mu}^{(1,1)} \rightarrow 2j)}{\Gamma(B_{\mu}^{(1,1)} \rightarrow 2t)}$
 - This ratio is *sensitive* to the cut-off scale M_s only through $B_{\mu}^{(1,1)}$ **mass**.

Continued...



- **Signals** are always well above the 5σ fluctuations of the **backgrounds**. This opens up a possibility, to measure the relative strengths of the signals in $2j$ and $2t$ channel.
 - One can determine $\frac{\Gamma(B_\mu^{(1,1)} \rightarrow 2j)}{\Gamma(B_\mu^{(1,1)} \rightarrow 2t)}$
 - This ratio is *sensitive* to the cut-off scale M_s only through $B_\mu^{(1,1)}$ **mass**.
 - **Mass** can be measured independently from the **peak position of E_γ distribution**.

Continued...



- **Signals** are always well above the 5σ fluctuations of the **backgrounds**. This opens up a possibility, to measure the relative strengths of the signals in $2j$ and $2t$ channel.
 - One can determine $\frac{\Gamma(B_{\mu}^{(1,1)} \rightarrow 2j)}{\Gamma(B_{\mu}^{(1,1)} \rightarrow 2t)}$
 - This ratio is *sensitive* to the cut-off scale M_s only through $B_{\mu}^{(1,1)}$ **mass**.
 - **Mass** can be measured independently from the **peak position of E_{γ} distribution**.
 - **Theoretical** number can be compared with the **experimentally** measured ratio of **decay widths**.

Results

Number of $\gamma + 2j$ **signal** and **SM background** events.
(assuming **500** fb^{-1} integrated luminosity).

e^+e^- C-o-M Energy	R^{-1} in GeV	$B_\mu^{(1,1)}$			$W_\mu^{3(1,1)}$		
		$m_{B_\mu^{(1,1)}}$ GeV	Signal Event	Background Event	$m_{W_\mu^{3(1,1)}}$ GeV	Signal Event	Background Event
500 GeV	280	387.3	5900	19258 (139)	433.8	253	26593 (163)
	290	401.1	6713	20368 (143)	448.7	349	34031 (184)
	300	414.9	7701	22207 (149)	463.7	520	50011 (224)
	310	428.8	9005	24814 (158)	478.7	-	-
	340	470.3	24296	59938 (245)	523.6	-	-
1 TeV	300	414.9	348	2889 (54)	463.7	10	2499 (50)
	400	553.3	430	2038 (45)	613.8	14	1932 (44)
	550	760.8	948	2096 (46)	840.4	43	2538 (50)
	630	871.4	2082	3013 (55)	961.5	210	8444 (92)
	690	954.4	6552	7482 (87)	1052.4	-	-

Concluding Remarks

- **2UED**: No. of **Extra Dimension** is 2, flat and compactified.
- **Spinless Adjoints** are the characteristics of this theory.
- If kinematically possible **Spinless Adjoints** dominantly decay to $t\bar{t}$, otherwise $b\bar{b}$ is not the dominant decay mode.
- Pair poroduction of coloured **Spinless Adjoints** vary from pb to few fb.
- Single Production of **EW Spinless Adjoints** at the **LHC** are miniscule.
- Discovery of $B_\mu^{(1,1)}$ in association with a **photon** at the **ILC** is possible for all R^{-1} , if kinematically possible.